

Numerical Ocean Models

- 1-2 km resolution models in Monterey Bay and Southern CA Bight
- 3 km resolution model covering ocean off all of California
- Drop-a-drifter and other trajectory tools
- Cross-sections
- 12 km resolution model running for area off whole West Coast




Jan. 8, 2013



1-2 km resolution models in Monterey Bay and Southern CA Bight

<http://www.cencoos.org/sections/models/ROMS.shtml>


Temp (°C, color), Current (m/s, arrows) at 0m for 12/17/2012 at 3GMT



<http://www.sccoos.org/data/roms/>

Regional Ocean Model System (ROMS) Model Output

The ROMS model is produced and distributed by NASA JPL and is available from <http://ocean.jpl.nasa.gov/SCB/>.




UCLA/JPL ROMS

Monterey Bay

3

- 1.5 km resolution
- T, S, velocity, SSH
- Nowcast every 6 hr
- 48 hr forecast
- NetCDF data files available

Temp (°C, color), Current (m/s, arrows) at 0m for 04/14/2012 at 3GM



"Drop-a-Drifter" tool

<http://ocean.jpl.nasa.gov/MB/mbmangen.jsp>

"Drop-a-Drifter" Surface Water Trajectories in Central California

Click the image below to view this Google Map based tool at the website for **NASA's Jet Propulsion Lab (JPL)** where the product was created. The movement of a particle of surface water (the trajectory) can be computed during the past (since Oct. 4th, 2010), present and future (48-hrs ahead). This can be used to indicate where any floating substance (oil etc.) or object (buoy etc.) came from or will go. Choose a starting point for a trajectory by "dropping" a virtual drifter marker on the map. The drifter path will be indicated by a colored line based on the dates selected. If you wish to determine the origin of something found floating or on the beach, please use the multiple drop mode and: select an end time of whenever it was found and a start time previous to the end time, then drop multiple drifters over a large area to see the most likely origin. Trajectories are determined using the **Regional Ocean Modeling System (ROMS)**. Note: although several data sources are incorporated in ROMS, the drifter tool has not yet been validated with actual data.



Get to this under Data Products menu

For multiple drifters,

Click to see push pin;

Drag mouse to get box

Click again

Choose compute & plot

Drop-a-Drifter Surface Water Trajectories Product

"Drop-a-Drifter" Surface Water Trajectories

http://www.cencoos.org/sections/products/drop_a_drifter.shtml

<http://ocean.jpl.nasa.gov/SCB/scbmangen.jsp>

If you wish to determine the origin of something found floating or on the beach, please use the multiple drop mode and: select an end time of whenever it was found and a start time previous to the end time, then drop multiple drifters over a large area to see the most likely origin.

☐ Single Drop Mode ☒ Multiple Drop Mode
(Click & then move the mouse to draw a region for drifters)

Number Of Drifters:

X dimension: Y dimension:

Location

Lon: Lat:

Data Source

ROMS 72hr Forecast


Start Time (GMT)

2012-09-06 03:00:00

End Time (GMT)

2012-09-09 03:00:00



Parameters



3 km resolution model covering ocean off all of CA⁶



<http://ourocean.jpl.nasa.gov/CA/>

Temp (°C, color), Current (m/s, arrows) at 0m for 01/01/2013




Temperature
and velocity

Temp (°C, color), Current (m/s, arrows) at 0m for 01/01/2013 at 3GM



UCLA/JPL ROMS - California

- 3 km resolution
- T, S, velocity, SSH
- Nowcast every 6 hr
- 72 hr forecast
- Data files available via THREDDS




Through the SCCOOS interface to the CA ROMS

www.sccoos.org/data/roms-3km


- Different depths, times, and variables, can be chosen for display
- Vertical profiles at a point may be plotted
- Cross-sections of different variables may be plotted

Note: does not work with Internet Explorer



CeNCOOS product under development


- Based on CA ROMS 72-hr current forecasts
- “Drifters” start at every 10th model grid point
- 1/2 hr time steps are used for the advection; black “tail” is 12 time steps (6 hrs) long
- Gray tail extends back to the starting location.
- Temperature field is updated every hour
- Temperature contours are every degree from 6 to 20 °C



12 km resolution model for whole West Coast

http://www.cencoos.org/sections/models/UCSC_ROMS.shtml

- 4D VAR
- 1/10 deg resolution
- T, S, velocity, SSH
- Figures updated daily
- Hindcast and nowcast
- NetCDF data files available
- Developed and run by UCSC



Biological and geochemical forecast models under development or consideration

- Statistical models relating ocean conditions to HABs
- Linked hydrologic, ocean, atmosphere models to forecast salmon populations
- Coupled physical/NPZD (nitrogen, phytoplankton, zooplankton, detritus) models for ecosystem forecasts
- Geochemical modeling to identify natural and outfall-based sources of nutrients




Jan. 8, 2013



Geoportal viewer integrating multiple asset types

e.g. <http://assets.maracoos.org>



Case Study 1: Oil Spills

- Cosco Busan hit Oakland Bay Bridge Nov. 7, 2007
 - 2004-present Establishment of California HF Radar array
 - 2005 IOOS office supports CORDC to develop HFR national network distribution
 - 2006 SafeSeas06 was the first collaborative effort between OOS and NOAA OR&R netCDF format standardized,
 - 2007 *CoscoBusan* incident used HFR in spill predictions, development of GIS format
 - 2008 HF Radar integration into both Office of Spill Prevention and Response (OSPR) and Office of Response and Restoration for GNOME forecasting model and used in National Preparedness for Emergency Response Planning (NPREP)
- Deepwater Horizon well blowout Apr. 20, 2010
 - April 20, 2010 No Gulf of Mexico HF Radar systems online
 - April 24, 2010 University of Southern Florida Systems online in national network
 - May 01, 2010 University of Southern Mississippi systems online in national network





Jan. 8, 2013



Cosco Busan, San Francisco Bay



Gap filled data, Trajectories




Oil spilled or flowing into offshore waters is within an environment where HF radar observations can be uniquely helpful in tracking and predicting movements

CeNCOOS and Cosco Busan Spill

http://www.cencoos.org/sections/news/SF_oil_spill_2007.shtml





Predicted Drifter Tracks for 10 November 2007 GMT

Based on Predicted Surface Currents

- Predicted Drifter Track
- Track Starting Point
- ⚡ HF Radar Antenna Locations

Farallon Islands

San Francisco

**GIS format
requested
by OSPR**

Data Courtesy of COCMP and CeNCOOS


Not Intended for Navigation



Working Draft
Do Not Circulate

Tim Reed, SIMoN/GFNMS, 10 Nov 2007

Oil spill response: OSPR-generated response exercise map with integrated HFR surface current data (arrows show speed and direction)

18






Improved network
response and data
integration between
Cosco Busan and
Deepwater Horizon

Mapping Surface Currents in Gulf of Mexico with HF Radar

Scripps Point of Contact: Eric Terrill (eterrill@ucsd.edu)
www.cordc.ucsd.edu/projects/mapping



Scripps HF Radar Activities in Gulf Of Mexico

- Developers of real-time data management and distribution system for national network (120+ radars). Leverages State of CA network.
- Processing of data for NOAA OR&R for official trajectories.
- Real-time Google products for data overlays.
- BP funded project to develop capability for HF radar on oil platforms

SAR imagery from CSTARS/UMIAMI. OI mapped HF Radar surface currents from CORDC/SIO. Radars operated by USM.


©2010 Europa Technologies
Data SIO, NOAA, U.S. Navy, NGA, GEBCO
28°42'57"N 88°5'74"W elev -5886 ft

Eye alt 352.97 mi


Integrated observations and model forecasts

21

<http://rucool.marine.rutgers.edu/deepwater>




Subsurface Monitoring Using Autonomous Vehicles



June 7 - July 26, 2010

Ocean Currents



Case Study 2: Outfall and Coastal Plume Tracking


- SF Ocean Beach Outfall Accidental Release
- Orange County Sanitation District Outfall Diversion (September 2012)



Jan. 8, 2013




SF Sewage District Ocean Beach Outfall Discharge October, 2007



2012 Orange County Sanitation District (OCSD)²⁵ Ocean Outfall Diversion

The Orange County Sanitation District (OCSD) discharges its treated effluent from a 120-inch ocean outfall that terminates in 200 feet of water, approximately 4.5 miles offshore Newport Beach and Huntington Beach.

On September 14, 2012 OCSD diverted the flow from the 120-inch outfall to the 78-inch outfall as part of a project to inspect, assess, and rehabilitate the Outfall Land Section and Ocean Outfall Booster Pump Station Piping.





The District has a secondary, 78-inch outfall located in about 60 feet of water, 1 mile off the coast. Periodically, OCSD request special permits to divert effluent from the 120-inch pipe to the 78-inch pipe for emergency purposes and planned maintenance projects.

2012 OCSD Ocean Outfall Diversion

<http://www.sccoos.org/projects/ocsd-diversion/>


Asset Map and
Visualization
Mooring, gliders,
HF radar, casts,
Samples





Case Study 3: Tar Balls

Late January 2008,
oil and tarballs
washed up on
beaches from
Monterey to San
Francisco


They were
determined to be
from a natural seep,
not from the Cosco
Busan or other spil



Surface current measurements showed that the tarballs' appearance followed a period of exceptionally strong flow from the south



Northward flow continuous from Santa Barbara to San Francisco




Hypoxia

- Low oxygen events have caused massive fish kills in harbors, like the one in Redondo Beach in March 2011
- Large scale low oxygen areas (also called anoxic or dead zones) have been observed off WA and OR. These have expanded from the continental shelf to near-shore waters every summer since 2002. The close proximity of these dead zones to the shore had never been reported before that year.




Low-oxygen conditions (<1.4 ml/l, blue) and extremely low-oxygen conditions (<0.5 ml/l, purple)




Ocean Acidification Efforts

It is important to note that OA efforts are in a preliminary stage



OA related time series along the west coast from NOAA Observing workshop in June 2012



Much of current carbon chemistry along the US west coast are unsuitable for needs of C-CAN stakeholders

Ocean Acidification Focus

<http://www.pmel.noaa.gov/co2/OA2012Workshop/>
<http://c-can.msi.ucsb.edu/>

- Raise awareness: OOSes can connect community to academics for expertise
- Shore Station Sensors – initially
- ID & test available technology for suitability
- Document best practices for instrumentation
- Integrate data handling and dissemination
- Provide education and training for participants